

OUTER PLETHYSM, BURNSIDE RING AND β -RINGS

Gustavo OCHOA

Departamento de Matemáticas, Universidad de Zaragoza, 50009 Zaragoza, Spain

Communicated by K.W. Gruenberg

Received 25 April 1986

Revised 5 May 1987

Consider the graded ring $B(S) = \bigoplus_{n \geq 0} B(\Sigma_n)$, where $B(\Sigma_n)$ is the Burnside ring of the symmetric group of degree n , Σ_n . Our main aim is to provide $B(S)$ with the structure of operator ring by introducing a third operation, ' \vee ', called outer plethysm, which satisfies the suitable identities. Also, we establish the basic ideas for the study of a β -rings theory.

1. Introduction

The set \mathcal{A} of all λ -operations (natural transformations from the underlying set functor, from the category of λ -rings to the category of sets, to itself) is a λ -ring [10, Chapter I].

The free λ -ring in one generator, $Z[u_1, u_2, \dots]$ (where $\lambda^k(u_1) = u_k$, $\forall k \geq 1$), and the graded ring $R(S) = \bigoplus_{n \geq 0} R(\Sigma_n)$ (where $R(\Sigma_n)$ is the representation ring of the symmetric group of degree n , Σ_n) are λ -rings isomorphic to \mathcal{A} [1, 9, 10].

Knutson conjectures in [10] that the operation of $R(S)$ (called outer plethysm) which corresponds with the composition of \mathcal{A} can be described through a suitable wreath product. Hoffman proves and describes this connection in [9].

Now, consider the abelian group $B(S) = \bigoplus_{n \geq 0} B(\Sigma_n)$, where $B(\Sigma_n)$ is the Burnside ring of the group Σ_n (see [6]); then, $B(S)$ becomes a graded ring with the cross product, ' \times ', defined in Section 4.

In this paper we provide $B(S)$ with the structure of *operator ring* by defining a third operation, ' \vee ' (outer plethysm), which satisfies the identities of Definition 4.4. Observe that ' \vee ' will not be right additive, therefore we have to define ' $b \vee a$ ' for every a in $B(S)$ and not merely for an ' a ' in some $B(\Sigma_n)$. We use only algebraic techniques, avoiding the topological description of the Burnside ring given by Dieck [4].

Similar results to Lemmas 4.7 and 4.9 can be found in Brennan's thesis [13, Chapter III], and they have been included here in order to make this paper more selfcontained.

Finally, in Section 5 we point out some notes about β -rings.

Notation

In general, we follow the notation of [9], but we write $R(S)$ instead of S .

If $\psi: \Omega \rightarrow \Gamma$ is a group homomorphism, ψ^* (resp. ψ_*) will be the restriction (resp. induction) from $R(\Gamma)$ or $B(\Gamma)$ to $R(\Omega)$ or $B(\Omega)$ (resp. from $R(\Omega)$ or $B(\Omega)$ to $R(\Gamma)$ or $B(\Gamma)$).

The existence and properties of restriction and induction maps in the Burnside ring of a finite group are recalled in Section 3.

The wreath product $\Gamma \sim \Sigma_k$ (natural action) will be denoted by $\Sigma_k \langle \Gamma \rangle$. Also, we use the symbol ' $\underline{\times}$ ' to denote the cross product instead of ' \times '.

\mathbb{N} is the set of nonnegative integers. For each k in \mathbb{N} , an r -partition of k is a sequence (i_1, \dots, i_r) such that i_j is in \mathbb{N} for $j=1, \dots, r$ and $i_1 + \dots + i_r = k$.

In Section 2 we identify the rings $R(\Gamma_1 \times \Gamma_2)$ and $R(\Gamma_1) \otimes R(\Gamma_2)$, for all finite groups Γ_1 and Γ_2 .

2. The operator ring $R(S)$

As we have already recalled, the operation 'composition' of \mathcal{A} is transferred to $R(S)$ through the isomorphism $\mathcal{A} \simeq R(S)$, and then $R(S)$ becomes an operator ring. Hoffman [9, p. 28–29] proves that the behaviour of this operation ' \vee ' for elements $b \in R(\Sigma_k)$, $a \in R(\Sigma_l)$ is

$$b \vee a = \theta_*[(\bar{\tau}_k(a))(\pi^*(b))]$$

where the group homomorphisms

$$\theta: \Sigma_k \langle \Sigma_l \rangle \rightarrow \Sigma_{kl}, \quad \pi: \Sigma_k \langle \Sigma_l \rangle \rightarrow \Sigma_k$$

are the inclusion and the projection, respectively, and the map

$$\bar{\tau}_k: R(\Sigma_l) \rightarrow R(\Sigma_k \langle \Sigma_l \rangle)$$

acts on actual representations by

$$\bar{\tau}_k(M) = M \otimes \dots \otimes M$$

with the obvious action of $\Sigma_k \langle \Sigma_l \rangle$ (see [9, p. 12, ...]).

Up to now, ' $b \vee a$ ' has not been described when $a = a_1 + \dots + a_r$ with a_j in $R(\Sigma_{l_j})$ for $j=1, \dots, r$ (there is no problem concerning b due to the left additivity of ' \vee '). In order to suggest the 'good' definition of the operation ' \vee ' in $B(S)$ we give such a description, of which the proof is omitted.

Proposition 2.1. *Let $b \in R(\Sigma_k)$, $a_j \in R(\Sigma_{l_j})$ ($j=1, \dots, r$). Then*

$$b \vee (a_1 + \dots + a_r) = \sum_{\alpha} (\theta_{\alpha})_* \left[\left(\bigotimes_j \bar{\tau}_{l_j}(a_j) \right) (\pi_{\alpha}^* \delta_{\alpha}^*(b)) \right] \quad (1)$$

where the summation is over r -partitions $\alpha = (i_1, \dots, i_r)$ of k ; θ_α is the composition

$$\begin{aligned} \Sigma_{i_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_r} \langle \Sigma_{l_r} \rangle &\rightarrow \Sigma_{i_1 l_1} \times \cdots \times \Sigma_{i_r l_r} \rightarrow \Sigma_{i_1 l_1 + \cdots + i_r l_r}, \\ \bigotimes_j \bar{\tau}_{i_j}(a_j) &:= \bar{\tau}_{i_1}(a_1) \otimes \cdots \otimes \bar{\tau}_{i_r}(a_r) \in R(\Sigma_{i_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_r} \langle \Sigma_{l_r} \rangle), \\ \pi_\alpha : \Sigma_{i_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_r} \langle \Sigma_{l_r} \rangle &\rightarrow \Sigma_{i_1} \times \cdots \times \Sigma_{i_r}, \\ \delta_\alpha : \Sigma_{i_1} \times \cdots \times \Sigma_{i_r} &\rightarrow \Sigma_k. \quad \square \end{aligned}$$

3. Restriction and induction in the Burnside ring

If Γ is a finite group, by $B(\Gamma)$ we shall denote the Burnside ring of Γ as defined by Dress [6]; this is the Grothendieck ring of the semiring of the isomorphism classes of finite Γ -sets, where the sum is defined by the disjoint union and the product by the Cartesian product with diagonal action. If X is a Γ -set, we shall denote also by X the corresponding element in $B(\Gamma)$.

Induction 3.1. Let $\phi : \Gamma \rightarrow \Omega$ be a group homomorphism, X a Γ -set. In the Cartesian product $\Omega \times X$ we define the equivalence relation \sim given by $(g, x) \sim (g_1, x_1)$ if and only if there exists h in Γ such that $g_1 = g\phi(h^{-1})$ and $x_1 = hx$.

We shall denote the equivalence class of (g, x) by $(\overline{g, x})$.

The set $\Omega \times_\phi X = \Omega \times X / \sim$ is an Ω -set with the action

$$g_1(\overline{g, x}) = (\overline{g_1 g, x}), \quad \forall g_1, g \in \Omega, x \in X$$

and it is called the Ω -set induced from X by ϕ , and denoted by $\phi_*(X)$.

By additivity, ϕ_* is extended to a homomorphism of abelian groups

$$\phi_* : B(\Gamma) \rightarrow B(\Omega).$$

Restriction 3.2. Let $\phi : \Gamma \rightarrow \Omega$ be a group homomorphism and X an Ω -set. Then X becomes a Γ -set, denoted by $\phi^*(X) = X_\phi$, with respect to the action

$$hx := \phi(h)x, \quad \forall h \in \Gamma.$$

ϕ^* is extended to a homomorphism of abelian groups

$$\phi^* : B(\Omega) \rightarrow B(\Gamma).$$

Proposition 3.3. Let $\phi : \Gamma \rightarrow \Omega$ and $\psi : \Omega \rightarrow \Gamma$ be homomorphisms of finite groups. Then

- (i) ϕ^* is a homomorphism of commutative rings with unit;
- (ii) $(\psi\phi)^* = \phi^*\psi^*$;
- (iii) $(\psi\phi)_* = \psi_*\phi_*$;

- (iv) If ϕ is an isomorphism, then $\phi_*\phi^* = \text{id}_{B(\Omega)}$, $\phi^*\phi_* = \text{id}_{B(\Gamma)}$;
 (v) If $\Gamma = \Omega$ and ϕ is an inner automorphism, then $\phi^* = \phi_* = \text{id}_{B(\Gamma)}$. \square

Theorem 3.4 (Mackey). *If $\alpha: \Gamma \rightarrow \Lambda$ and $\beta: \Omega \rightarrow \Lambda$ are inclusions of subgroups, then*

$$\beta^*\alpha_* = \sum_g (\beta_g)^*(\alpha_g)_*$$

where g ranges over a set of (Γ, Ω) -double coset representatives, $\alpha_g: \Gamma^g \cap \Omega \rightarrow \Gamma$ is given by $\alpha_g(x) = gxg^{-1}$, and $\beta_g: \Gamma^g \cap \Omega \rightarrow \Omega$ is the inclusion. \square

Theorem 3.5 (Frobenius). *Let $\alpha: \Gamma \rightarrow \Omega$ be a group homomorphism. For every x in $B(\Gamma)$ and every y in $B(\Omega)$, one has*

$$\alpha_*(x) \cdot y = \alpha_*(x \cdot \alpha^*(y)). \quad \square$$

The proofs of Proposition 3.3 and Theorems 3.4 and 3.5 are standard. Dress comments the existence of these properties in [7, p. 42] in connection with the idea of $B(-)$ as G -functor [8].

Another property that we will need is the following:

Proposition 3.6. *If the diagram is a pullback of finite groups, then $\pi_1^*\alpha_* = \beta_*\pi^*$.*

$$\begin{array}{ccc} \Omega & \xrightarrow{\beta} & \Omega_1 \\ \pi \downarrow & & \downarrow \pi_1 \\ \Gamma & \xrightarrow{\alpha} & \Gamma_1 \end{array} \quad \square$$

The proof is a routine checking.

4. Operator ring structure in $B(S)$

As we have already mentioned, $B(S) = \bigoplus_{k \geq 0} B(\Sigma_k)$ is an abelian group. The following ring homomorphism will be very useful:

Definition 4.1. If Γ_1, Γ_2 are finite groups, and (for $i=1,2$) $\pi_i: \Gamma_1 \times \Gamma_2 \rightarrow \Gamma_i$ denote the projections, the homomorphism $\mu: B(\Gamma_1) \otimes B(\Gamma_2) \rightarrow B(\Gamma_1 \times \Gamma_2)$ is defined as the composition of $\pi_1^* \otimes \pi_2^*$ followed by the juxtaposition

$$B(\Gamma_1 \times \Gamma_2) \otimes B(\Gamma_1 \times \Gamma_2) \rightarrow B(\Gamma_1 \times \Gamma_2).$$

This can be done in the obvious way for more than two factors. Sometimes the notation $x \otimes y$ will be used instead of $\mu(x \otimes y)$.

Let Γ be a finite group and $P = \prod_{k \geq 0} B(\Sigma_k \langle \Gamma \rangle)$. The definition of the cross product, $\underline{\times}$, is (analogous to [9, p. 6]):

$$P \otimes P \xrightarrow{\underline{\times}} P$$

with components

$$B(\Sigma_i \langle \Gamma \rangle) \otimes B(\Sigma_j \langle \Gamma \rangle) \xrightarrow{\mu} B(\Sigma_i \langle \Gamma \rangle \times \Sigma_j \langle \Gamma \rangle) \xrightarrow{\phi_*} B(\Sigma_{i+j} \langle \Gamma \rangle)$$

where $\phi : \Sigma_i \langle \Gamma \rangle \times \Sigma_j \langle \Gamma \rangle \rightarrow \Sigma_{i+j} \langle \Gamma \rangle$ is the obvious group homomorphism.

With this multiplication $B(S)$ becomes a graded ring ($\Gamma=1$).

Also, we define the multiplication \otimes on the group $\bigoplus_{l=0}^{\infty} B(\Sigma_k \langle \Sigma_l \langle \Gamma \rangle \rangle)$ (see [9, p. 8]) given on components by the commutativity of the diagram

$$\begin{array}{ccc} B(\Sigma_k \langle \Sigma_i \langle \Gamma \rangle \rangle) \otimes B(\Sigma_k \langle \Sigma_j \langle \Gamma \rangle \rangle) & \xrightarrow{\otimes} & B(\Sigma_k \langle \Sigma_{i+j} \langle \Gamma \rangle \rangle) \\ \mu \downarrow & & \uparrow \alpha_* \\ B(\Sigma_k \langle \Sigma_i \langle \Gamma \rangle \rangle \times \Sigma_k \langle \Sigma_j \langle \Gamma \rangle \rangle) & \xrightarrow{\delta^*} & B(\Sigma_k \langle (\Sigma_i \times \Sigma_j) \langle \Gamma \rangle \rangle) \end{array}$$

where μ is given in Definition 4.1 and

$$\begin{aligned} \delta : \Sigma_k \langle (\Sigma_i \times \Sigma_j) \langle \Gamma \rangle \rangle &\rightarrow \Sigma_k \langle \Sigma_i \langle \Gamma \rangle \rangle \times \Sigma_k \langle \Sigma_j \langle \Gamma \rangle \rangle, \\ \alpha : \Sigma_k \langle (\Sigma_i \times \Sigma_j) \langle \Gamma \rangle \rangle &\rightarrow \Sigma_k \langle \Sigma_{i+j} \langle \Gamma \rangle \rangle \end{aligned}$$

are the obvious group homomorphisms.

Now, we define the map T from the set of isomorphism classes of simple Γ -sets to P , with components T_k given by

$$T_k(X) = X \times \dots \times X = X^k.$$

$T_k(X)$ is a $\Sigma_k \langle \Gamma \rangle$ -set with the action

$$(g_1, \dots, g_k; \sigma)(x_1, \dots, x_k) = (g_1 x_{\sigma^{-1}(1)}, \dots, g_k x_{\sigma^{-1}(k)}).$$

Proposition 4.2 (see [9, (3.7) and p. 112]). *T can be extended to a homomorphism from the additive group $B(S)$ to the group P with the cross product. (See [9, (1.6) and (2.3)]).*

Moreover, T satisfies the following properties:

- (o) For every Γ -set Y , $T_k(Y) = Y \times \dots \times Y$ with the same action as above;
- (i) $T_0(x) = 1$ and $T_1(x) = x$, for every $x \in B(\Gamma)$;
- (ii) $T(x + y) = T(x) \underline{\times} T(y)$, for every $x, y \in B(\Gamma)$;
- (iii) The following diagram commutes:

$$\begin{array}{ccc}
 B(\Gamma) & \xrightarrow{T^{\otimes r}} & P \otimes \cdots \otimes P \\
 T \downarrow & & \downarrow \bar{\mu} \\
 P & \xrightarrow{\Delta^{(r)}} & \prod B(\Sigma_{k_1}\langle\Gamma\rangle \times \cdots \times \Sigma_{k_r}\langle\Gamma\rangle)
 \end{array}$$

where the product ranges over all the $k_1, \dots, k_r \geq 0$; $\bar{\mu}$ has components

$$B(\Sigma_{k_1}\langle\Gamma\rangle) \otimes \cdots \otimes B(\Sigma_{k_r}\langle\Gamma\rangle) \xrightarrow{\mu} B(\Sigma_{k_1}\langle\Gamma\rangle \times \cdots \times \Sigma_{k_r}\langle\Gamma\rangle)$$

and $\Delta^{(r)}, \Delta_{k_1, \dots, k_r}^{(r)} = \alpha_{k_1, \dots, k_r}^*$ defined from

$$\alpha_{k_1, \dots, k_r} : \Sigma_{k_1}\langle\Gamma\rangle \times \cdots \times \Sigma_{k_r}\langle\Gamma\rangle \rightarrow \Sigma_k\langle\Gamma\rangle \quad (k = k_1 + \cdots + k_r);$$

(iv) $T(xy) = T(x)T(y)$ for every $x, y \in B(\Gamma)$;

(v) The following diagram commutes:

$$\begin{array}{ccc}
 B(\Gamma) & \xrightarrow{T} & P \\
 T \downarrow & & \downarrow \zeta \\
 P & \xrightarrow{\square} & \prod_{k, l \geq 0} B(\Sigma_k\langle\Sigma_l\langle\Gamma\rangle\rangle)
 \end{array}$$

where ζ has components

$$\zeta_{k, l} : B(\Sigma_l\langle\Gamma\rangle) \rightarrow B(\Sigma_k\langle\Sigma_l\langle\Gamma\rangle\rangle)$$

which act on $B(\Sigma_l\langle\Gamma\rangle)$ in the same way as $T_k(\Sigma_l\langle\Gamma\rangle)$ playing the role of Γ . We will sometimes denote $\zeta_{k, l} = T_k$.

\square has components $\square_{k, l} = \theta_{k, l}^*$, where

$$\theta_{k, l} : \Sigma_k\langle\Sigma_l\langle\Gamma\rangle\rangle \rightarrow \Sigma_{kl}\langle\Gamma\rangle$$

is defined by

$$\begin{aligned}
 &\theta_{k, l}[(g_{11}, \dots, g_{1l}; \sigma_1), \dots, (g_{k1}, \dots, g_{kl}; \sigma_k); \tau] \\
 &= [g_{11}, \dots, g_{1l}, \dots, g_{k1}, \dots, g_{kl}; (\sigma_1, \dots, \sigma_k; \tau)].
 \end{aligned}$$

Sketch of proof. (o), (i) and (ii) are not difficult.

(iii) It is enough to prove that both $\mu T^{\otimes r}$ and $\Delta^{(r)}T$ coincide on Γ -sets, their images are contained in the set of \times -invertibles of $B(\Sigma_{k_1}\langle\Gamma\rangle \times \cdots \times \Sigma_{k_r}\langle\Gamma\rangle)$, and are homomorphisms from '+' to ' \times '; then proceed in the same way as the proof of [9, (3.7) III].

(iv) Check the statement for X, Y Γ -sets and then follow the proof of [9, Proposition (3.7) IV].

(v) The diagram commutes for Γ -sets. Now, the proof is similar to [9, (3.7) V]. \square

Lemma 4.3. *Let $a \in B(\Sigma_i \langle \Gamma \rangle)$, $b \in B(\Sigma_j \langle \Gamma \rangle)$. Then*

$$T_k(a \times b) = T_k(a) \otimes T_k(b).$$

Proof. Follow step by step the proof of [9, Proposition (2.10)]. \square

Our main aim is to provide $B(S)$ with the structure of operator ring. Let us recall what this means:

Definition 4.4. An *operator ring* is a commutative ring R with identity 1, and with a chosen element $e (\neq 1)$ of R and a composition $\vee : R \times R \rightarrow R$ satisfying:

- (1) $(x_1 + x_2) \vee x_3 = (x_1 \vee x_3) + (x_2 \vee x_3)$,
- (2) $(x_1 x_2) \vee x_3 = (x_1 \vee x_3)(x_2 \vee x_3)$,
- (3) $(x_1 \vee x_2) \vee x_3 = x_1 \vee (x_2 \vee x_3)$,
- (4) $x_1 \vee e = x_1$,

for all x_1, x_2, x_3 in R .

We want to define the operation ‘ \vee ’ in $B(S)$. Let k, l_1, \dots, l_r be elements in \mathbb{N} . For every r -partition of k , $\alpha = (i_1, \dots, i_r)$, one has the group homomorphisms θ_α , δ_α and π_α mentioned in Proposition 2.1.

Definition 4.5. Let $b \in B(\Sigma_k)$, $a_j \in B(\Sigma_{l_j})$ ($j=1, \dots, r$). We define the composition ‘ \vee ’ in $B(S)$ by

$$b \vee (a_1 + \dots + a_r) = \sum_{\alpha} (\theta_{\alpha})_* \left[\left(\bigotimes_{j=1}^r T_{l_j}(a_j) \right) (\pi_{\alpha}^* \delta_{\alpha}^*(b)) \right]$$

where the sum ranges over all r -partitions of k : $\alpha = (i_1, \dots, i_r)$.

In some cases an element a in $B(S)$ will not be represented as a sum $a = a_1 + \dots + a_r$ with each summand a_j appearing in a different $B(\Sigma_{l_j})$; so we have to prove that this composition does not depend on the representation of a .

Proposition 4.6. *The operation composition ‘ \vee ’ of Definition 4.5 is well defined.*

Proof. Let $a_1 = c_1 + d_1 \in B(\Sigma_{l_1})$, $a_j \in B(\Sigma_{l_j})$ ($j=2, \dots, r$). Put $l_0 := l_1$, $a'_0 := c_1$, $a'_1 := d_1$, $a'_j := a_j$ ($j=2, \dots, r$). For every $(r+1)$ -partition of k , $\alpha' = (i'_0, \dots, i'_r)$, consider the r -partition of k , $\alpha = (i'_0 + i'_1, i'_2, \dots, i'_r) = (i_1, i_2, \dots, i_r)$.

Let $\phi_{\alpha'}$ be the group homomorphism

$$\Sigma_{i'_0} \langle \Sigma_{l_0} \rangle \times \Sigma_{i'_1} \langle \Sigma_{l_1} \rangle \times \dots \times \Sigma_{i'_r} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i'_0 + i'_1} \langle \Sigma_{l_1} \rangle \times \Sigma_{i'_2} \langle \Sigma_{l_2} \rangle \times \dots \times \Sigma_{i'_r} \langle \Sigma_{l_r} \rangle$$

and $\theta_{\alpha'}$, θ_{α} , $\delta_{\alpha'}$, δ_{α} , $\pi_{\alpha'}$, π_{α} as before. It is easy to check that

$$\theta_{\alpha'} = \theta_{\alpha} \phi_{\alpha'} \quad \text{and} \quad \delta_{\alpha'} \pi_{\alpha'} = \delta_{\alpha} \pi_{\alpha} \phi_{\alpha'}.$$

So, we have $(\theta_{\alpha'})_* = (\theta_{\alpha})_*(\phi_{\alpha'})_*$ and $\pi_{\alpha'}^* \delta_{\alpha'}^* = \phi_{\alpha'}^* \pi_{\alpha}^* \delta_{\alpha}^*$.

Now,

$$\begin{aligned}
 & b \vee (a'_0 + a'_1 + \cdots + a'_r) \\
 &= \sum_{\alpha'} (\theta_{\alpha'})_* \left[\left(\bigotimes_{j=0}^r T_{i'_j}(a'_j) \right) (\pi_{\alpha'}^* \delta_{\alpha'}^*(b)) \right] \\
 &= \sum_{\alpha'} (\theta_{\alpha})_*(\phi_{\alpha'})_* \left[\left(\bigotimes_{j=0}^r T_{i'_j}(a'_j) \right) (\phi_{\alpha'}^* \pi_{\alpha}^* \delta_{\alpha}^*(b)) \right] \\
 &\quad \text{by Frobenius reciprocity} \\
 &= \sum_{\alpha'} (\theta_{\alpha})_* \left[\left((\phi_{\alpha'})_* \left(\bigotimes_{j=0}^r T_{i'_j}(a'_j) \right) \right) (\pi_{\alpha}^* \delta_{\alpha}^*(b)) \right] \\
 &\quad \text{when } \alpha' \text{ ranges over all } (r+1)\text{-partitions of } k, \alpha \text{ ranges over all} \\
 &\quad r\text{-partitions of } k. \text{ Put } P(\alpha) \text{ for the set of } (r+1)\text{-partitions } \alpha' \text{ de-} \\
 &\quad \text{fining the same } \alpha \\
 &= \sum_{\alpha} (\theta_{\alpha})_* \left[\left[\sum_{P(\alpha)} (T_{i'_0}(a'_0) \otimes T_{i'_1}(a'_1)) \right] \otimes T_{i'_2}(a'_2) \otimes \cdots \otimes T_{i'_r}(a'_r) \right] (\pi_{\alpha}^* \delta_{\alpha}^*(b)) \\
 &= \sum_{\alpha} (\theta_{\alpha})_* [(T_{i_1}(a'_0 + a'_1) \otimes T_{i_2}(a'_2) \otimes \cdots \otimes T_{i_r}(a'_r)) (\pi_{\alpha}^* \delta_{\alpha}^*(b))] \\
 &= b \vee (a_1 + \cdots + a_r). \quad \square
 \end{aligned}$$

Lemma 4.7 (see [9, Theorem (1.2)] and [3, Chapter III]). Let $n = \varepsilon_1 + \cdots + \varepsilon_r = \omega_1 + \cdots + \omega_s$ with $\varepsilon_i \in \mathbb{N}$ ($i = 1, \dots, r$) and $\omega_j \in \mathbb{N}$ ($j = 1, \dots, s$). Let H, K be the subgroups of Σ_n :

$$H = \Sigma_{\varepsilon_1} \times \cdots \times \Sigma_{\varepsilon_r}, \quad K = \Sigma_{\omega_1} \times \cdots \times \Sigma_{\omega_s}.$$

Let M be the set of all \mathbb{N} -matrices $r \times s$ with row sums $(\varepsilon_1, \dots, \varepsilon_r)$ and column sums $(\omega_1, \dots, \omega_s)$. Let $C = \{HgK : g \in \Sigma_n\}$.

Then there exists a bijective map $\Psi : M \rightarrow C$ such that if $\Psi[A = (a_{ij})] = HgK$, then

$$H^g \cap K = (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}}) \leq \Sigma_n.$$

Proof. Omitted. \square

Notations for Proposition 4.8

$$k = k_1 + k_2.$$

$\varepsilon = (i_1, \dots, i_r)$ is an r -partition of k .

$\varepsilon_1 = (i_{11}, \dots, i_{1r})$ is an r -partition of k_1 .

$\varepsilon_2 = (i_{21}, \dots, i_{2r})$ is an r -partition of k_2 .

$$\phi : \Sigma_{k_1} \times \Sigma_{k_2} \rightarrow \Sigma_k.$$

$$\delta_{\varepsilon_1} : \Sigma_{i_{11}} \times \cdots \times \Sigma_{i_{1r}} \rightarrow \Sigma_{k_1}.$$

$$\delta_{\varepsilon_2} : \Sigma_{i_{21}} \times \cdots \times \Sigma_{i_{2r}} \rightarrow \Sigma_{k_2}.$$

$$\delta_{\varepsilon} : \Sigma_{i_1} \times \cdots \times \Sigma_{i_r} \rightarrow \Sigma_k.$$

$$\begin{aligned}
\pi_{\varepsilon_1} &: \Sigma_{i_{11}} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_{1r}} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i_{11}} \times \cdots \times \Sigma_{i_{1r}}. \\
\pi_{\varepsilon_2} &: \Sigma_{i_{21}} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_{2r}} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i_{21}} \times \cdots \times \Sigma_{i_{2r}}. \\
\pi_{\varepsilon} &: \Sigma_{i_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_r} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i_1} \times \cdots \times \Sigma_{i_r}. \\
\theta_{\varepsilon_1} &: \Sigma_{i_{11}} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_{1r}} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i_{11}l_1} \times \cdots \times \Sigma_{i_{1r}l_r} \rightarrow \Sigma_{i_{11}l_1 + \cdots + i_{1r}l_r}. \\
\theta_{\varepsilon_2} &: \Sigma_{i_{21}} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_{2r}} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i_{21}l_1} \times \cdots \times \Sigma_{i_{2r}l_r} \rightarrow \Sigma_{i_{21}l_1 + \cdots + i_{2r}l_r}. \\
\alpha &: (\Sigma_{i_{11}} \times \Sigma_{i_{21}}) \times \cdots \times (\Sigma_{i_{1r}} \times \Sigma_{i_{2r}}) \rightarrow \Sigma_{i_1} \times \cdots \times \Sigma_{i_r}. \\
\bar{\alpha} &: (\Sigma_{i_{11}} \times \Sigma_{i_{21}}) \langle \Sigma_{l_1} \rangle \times \cdots \times (\Sigma_{i_{1r}} \times \Sigma_{i_{2r}}) \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{i_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_r} \langle \Sigma_{l_r} \rangle. \\
\bar{\alpha}_j &: (\Sigma_{i_{1j}} \times \Sigma_{i_{2j}}) \langle \Sigma_{l_j} \rangle \rightarrow \Sigma_{i_j} \langle \Sigma_{l_j} \rangle \quad (j=1, \dots, r). \\
\bar{\pi} &: (\Sigma_{i_{11}} \times \Sigma_{i_{21}}) \langle \Sigma_{l_1} \rangle \times \cdots \times (\Sigma_{i_{1r}} \times \Sigma_{i_{2r}}) \langle \Sigma_{l_r} \rangle \rightarrow (\Sigma_{i_{11}} \times \Sigma_{i_{21}}) \times \cdots \times (\Sigma_{i_{1r}} \times \Sigma_{i_{2r}}). \\
\gamma &: (\Sigma_{i_{11}} \times \Sigma_{i_{21}}) \times \cdots \times (\Sigma_{i_{1r}} \times \Sigma_{i_{2r}}) \rightarrow (\Sigma_{i_{11}} \times \cdots \times \Sigma_{i_{1r}}) \times (\Sigma_{i_{21}} \times \cdots \times \Sigma_{i_{2r}}). \\
\bar{\gamma} &: (\Sigma_{i_{11}} \times \Sigma_{i_{21}}) \langle \Sigma_{l_1} \rangle \times \cdots \times (\Sigma_{i_{1r}} \times \Sigma_{i_{2r}}) \langle \Sigma_{l_r} \rangle \\
&\quad \rightarrow (\Sigma_{i_{11}} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_{1r}} \langle \Sigma_{l_r} \rangle) \times (\Sigma_{i_{21}} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{i_{2r}} \langle \Sigma_{l_r} \rangle). \\
\psi &: \Sigma_{i_{11}l_1 + \cdots + i_{1r}l_r} \times \Sigma_{i_{21}l_1 + \cdots + i_{2r}l_r} \rightarrow \Sigma_{i_1l_1 + \cdots + i_rl_r}. \\
\sigma &: \Sigma_{i_1l_1 + \cdots + i_rl_r} \rightarrow \Sigma_{i_1l_1 + \cdots + i_rl_r} \text{ is a certain inner automorphism.}
\end{aligned}$$

Proposition 4.8. *Property 2 of Definition 4.4 is valid in $B(S)$.*

Proof. We have to prove $(b_1 \times b_2) \vee a = (b_1 \vee a) \times (b_2 \vee a)$ for all b_1, b_2, a in $B(S)$. By bi-additivity of \times and left additivity of \vee we may assume $b_1 \in B(\Sigma_{k_1}), b_2 \in B(\Sigma_{k_2})$.

Let $a = a_1 + \cdots + a_r$ with $a_j \in B(\Sigma_{l_j})$ ($j=1, \dots, r$).

$$\begin{aligned}
(b_1 \times b_2) \vee a &= \phi_*(b_1 \otimes b_2) \vee a \\
&= \sum_{\varepsilon} (\theta_{\varepsilon})_* \left[\left(\bigotimes_j T_{l_j}(a_j) \right) (\pi_{\varepsilon}^* \delta_{\varepsilon}^*(\phi_*(b_1 \otimes b_2))) \right] = \\
&\quad \text{by the Mackey theorem and Lemma 4.7} \\
&= \sum_{\varepsilon} (\theta_{\varepsilon})_* \left[\left(\bigotimes_j T_{l_j}(a_j) \right) \left(\pi_{\varepsilon}^* \left[\sum_{\varepsilon_1 + \varepsilon_2 = \varepsilon} \alpha_* \gamma^*(\delta_{\varepsilon_1} \otimes \delta_{\varepsilon_2})^*(b_1 \otimes b_2) \right] \right) \right] \\
&\quad \text{by applying Proposition 3.6 to } \pi_{\varepsilon} \bar{\alpha} = \alpha \bar{\pi} \text{ and putting } \varepsilon := \varepsilon_1 + \varepsilon_2 \\
&= \sum_{\varepsilon_1, \varepsilon_2} (\theta_{\varepsilon})_* \left[\left(\bigotimes_j T_{l_j}(a_j) \right) (\bar{\alpha}_* \bar{\pi}^* \gamma^*(\delta_{\varepsilon_1}^*(b_1) \otimes \delta_{\varepsilon_2}^*(b_2))) \right] \\
&\quad \text{since } \gamma \bar{\pi} = (\pi_{\varepsilon_1} \otimes \pi_{\varepsilon_2}) \bar{\gamma} \\
&= \sum_{\varepsilon} (\theta_{\varepsilon})_* \left[\left(\bigotimes_j T_{l_j}(a_j) \right) (\bar{\alpha}_* \bar{\gamma}^* [\pi_{\varepsilon_1}^* \delta_{\varepsilon_1}^*(b_1) \otimes \pi_{\varepsilon_2}^* \delta_{\varepsilon_2}^*(b_2)]) \right] = \\
&\quad \text{Now, } \bar{\alpha}^*(\bigotimes_j T_{l_j}(a_j)) = \bigotimes_j \bar{\alpha}_j(T_{l_j}(a_j)) = \bigotimes_j (T_{i_{1j}}(a_j) \otimes T_{i_{2j}}(a_j)) \text{ (the} \\
&\quad \text{last equality is due to Proposition 4.2(iii)). By Frobenius reciprocity} \\
&= \sum_{\varepsilon} (\theta_{\varepsilon})_* \bar{\alpha}_* \left[\left(\bigotimes_j [T_{i_{1j}}(a_j) \otimes T_{i_{2j}}(a_j)] \right) (\bar{\gamma}^* [\pi_{\varepsilon_1}^* \delta_{\varepsilon_1}^*(b_1) \otimes \pi_{\varepsilon_2}^* \delta_{\varepsilon_2}^*(b_2)]) \right] \\
&\quad \text{since } \theta_{\varepsilon} \bar{\alpha} = \sigma \psi(\theta_{\varepsilon_1} \otimes \theta_{\varepsilon_2}) \bar{\gamma} \text{ and by Frobenius reciprocity}
\end{aligned}$$

$$\begin{aligned}
&= \sum \sigma_* \psi_*(\theta_{\varepsilon_1} \otimes \theta_{\varepsilon_2})_* \left[\bar{\gamma}_* \left(\bigotimes_j [T_{i_{1j}}(a_j) \otimes T_{i_{2j}}(a_j)] \right) (\pi_{\varepsilon_1}^* \delta_{\varepsilon_1}^*(b_1) \otimes \pi_{\varepsilon_2}^* \delta_{\varepsilon_2}^*(b_2)) \right] \\
&\quad \text{since } \sigma_* = 1 \text{ by Proposition 3.3(v)} \\
&= \sum \psi_*(\theta_{\varepsilon_1} \otimes \theta_{\varepsilon_2})_* \left[\left(\left[\bigotimes_j T_{i_{1j}}(a_j) \right] \right. \right. \\
&\quad \left. \left. \otimes \left[\bigotimes_j T_{i_{2j}}(a_j) \right] \right) (\pi_{\varepsilon_1}^* \delta_{\varepsilon_1}^*(b_1) \otimes \pi_{\varepsilon_2}^* \delta_{\varepsilon_2}^*(b_2)) \right] \\
&= \sum \psi_* \left[(\theta_{\varepsilon_1})_* \left(\left[\bigotimes_j T_{i_{1j}}(a_j) \right] [\pi_{\varepsilon_1}^* \delta_{\varepsilon_1}^*(b_1)] \right) \right. \\
&\quad \left. \otimes (\theta_{\varepsilon_2})_* \left(\left[\bigotimes_j T_{i_{2j}}(a_j) \right] [\pi_{\varepsilon_2}^* \delta_{\varepsilon_2}^*(b_2)] \right) \right] \\
&= \left(\sum_{\varepsilon_1} (\theta_{\varepsilon_1})_* \left[\left(\bigotimes_j T_{i_{1j}}(a_j) \right) (\pi_{\varepsilon_1}^* \delta_{\varepsilon_1}^*(b_1)) \right] \right) \\
&\quad \times \left(\sum_{\varepsilon_2} (\theta_{\varepsilon_2})_* \left[\left(\bigotimes_j T_{i_{2j}}(a_j) \right) (\pi_{\varepsilon_2}^* \delta_{\varepsilon_2}^*(b_2)) \right] \right) \\
&= (b_1 \vee a) \times (b_2 \vee a). \quad \square
\end{aligned}$$

Lemma 4.9. Let $H = \Sigma_{\varepsilon_1} \langle \Sigma_{l_1} \rangle$ and $K = \Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{1s}}$, with $a_{11} + \cdots + a_{1s} = \varepsilon_1 l_1$, be subgroups of $\Sigma_{\varepsilon_1 l_1}$.

In the set of all \mathbb{N} -matrices $\varepsilon_1 \times s$ with row sums (l_1, \dots, l_1) and column sums (a_{11}, \dots, a_{1s}) define the equivalence relation \sim by $B_{\varepsilon_1} \sim B'_{\varepsilon_1}$ if and only if there exists a certain permutation of the ε_1 rows of B_{ε_1} to obtain B'_{ε_1} . The equivalence class of B_{ε_1} is written $[B_{\varepsilon_1}]$.

Let M be the set of all equivalence classes and $C = \{HgK : g \in \Sigma_{\varepsilon_1 l_1}\}$. Then there exists a bijective map

$$\Psi : M \rightarrow C$$

such that, if $\Psi([B_{\varepsilon_1}]) = HgK$ with $B_{\varepsilon_1} = (b_{ij})$, then $H \cap K^{g^{-1}} = \Sigma_{\varepsilon_1} \langle \Sigma_{\gamma(1)_1^1} \times \cdots \times \Sigma_{\gamma(1)_s^1} \rangle \times \cdots \times \Sigma_{e_{p_1}} \langle \Sigma_{\gamma(1)_1^{p_1}} \times \cdots \times \Sigma_{\gamma(1)_s^{p_1}} \rangle$ where $\{\gamma(1)_1^1, \dots, \gamma(1)_s^{p_1}\}$ is the set of all s -partitions of l_1 : $\gamma(1)^i = (\gamma(1)_1^i, \dots, \gamma(1)_s^i)$ ($i = 1, \dots, p_1$), and e_i is the number of rows of B_{ε_1} which are equal to $\gamma(1)^i$.

Proof. Let $[B_{\varepsilon_1}] \in M$, $B_{\varepsilon_1} = (b_{ij})$. We define $\Psi([B_{\varepsilon_1}]) = HgK$ with g given by:

For $l = (i-1)l_1 + b_{i1} + \cdots + b_{i(j-1)} + t$ ($1 \leq t \leq b_{ij}$)

$$lg = a_{11} + \cdots + a_{1(j-1)} + b_{1j} + \cdots + b_{(i-1)j} + t. \quad (2)$$

If $[\bar{B}_{\varepsilon_1}] = [B_{\varepsilon_1}]$ with $\bar{B}_{\varepsilon_1} = (\bar{b}_{ij})$, let $\sigma \in \Sigma_{\varepsilon_1}$ be such that

$$\bar{b}_{i\sigma, j} = b_{ij} \quad (1 \leq i \leq \varepsilon_1; 1 \leq j \leq s).$$

Also, we put $\sigma = (1, \dots, 1; \sigma) \in \Sigma_{\varepsilon_1} \langle \Sigma_{l_1} \rangle$.

Let l be as before and \bar{g} defined for \bar{B}_{e_1} as g for B_{e_1} . Now,

$$\begin{aligned} l\sigma &= (i\sigma - 1)l_1 + \bar{b}_{i\sigma,1} + \cdots + \bar{b}_{i\sigma,j-1} + t \quad (1 \leq t \leq \bar{b}_{i\sigma,j}), \\ (l\sigma)\bar{g} &= a_{11} + \cdots + a_{1,j-1} + \bar{b}_{1j} + \cdots + \bar{b}_{i\sigma-1,j} + t. \end{aligned}$$

So, $\sigma\bar{g} = g\tau$ for some $\tau \in K$. Then Ψ is well defined.

We define $\Phi: C \rightarrow M$ by $\Phi(HgK) = [B_{e_1}]$ with $B_{e_1} = (b_{ij})$ and b_{ij} the cardinal of the intersection

$$\{(i-1)l_1 + 1, \dots, il_1\} g \cap \{a_{11} + \cdots + a_{1,j-1} + 1, \dots, a_{11} + \cdots + a_{1j}\}.$$

Φ is well defined; in fact:

Let $\bar{g} = h g k$ ($h \in H$, $k \in K$) and suppose $h = (x_1, \dots, x_{e_1}; \sigma)$. Then \bar{g} yields the matrix $\bar{B}_{e_1} = (\bar{b}_{ij})$ with \bar{b}_{ij} the cardinal of each intersection:

$$\begin{aligned} &\{(i-1)l_1 + 1, \dots, il_1\} h g k \cap \{a_{11} + \cdots + a_{1,j-1} + 1, \dots, a_{11} + \cdots + a_{1j}\}, \\ &\{(i-1)l_1 + 1, \dots, il_1\} \sigma g \cap \{a_{11} + \cdots + a_{1,j-1} + 1, \dots, a_{11} + \cdots + a_{1j}\} k^{-1}, \\ &\{(i\sigma - 1)l_1 + 1, \dots, (i\sigma)l_1\} g \cap \{a_{11} + \cdots + a_{1,j-1} + 1, \dots, a_{11} + \cdots + a_{1j}\}. \end{aligned}$$

So, $\bar{b}_{ij} = b_{i\sigma,j}$ and then $[B_{e_1}] = [\bar{B}_{e_1}]$.

Clearly, $\Phi\Psi([B_{e_1}]) = [B_{e_1}]$.

When $\Phi(HgK) = [B_{e_1}]$ we may choose the representative g acting exactly as in (2) and then it is clear that

$$\Psi\Phi(HgK) = HgK.$$

Finally, suppose $\Psi([B_{e_1}]) = HgK$. Let us choose B_{e_1} such that the first e_1 rows are equal to $\gamma(1)^1$, the following e_2 rows are $\gamma(1)^2$, etc. The representative g acts as in (2). Then, $K^{g^{-1}}$ is the subgroup of $\Sigma_{e_1 l_1}$ of all permutations fixing the sets X_r for $1 \leq r \leq s$ where j is in X_r if and only if

$$1 + kl_1 + \sum_{i=1}^{r-1} b_{k+1,i} \leq j \leq kl_1 + \sum_{j=1}^r b_{k+1,i}$$

for some k with $0 \leq k < e_1$.

The elements in H are those of $\Sigma_{l_1 e_1}$ which permute the sets $\{1, 2, \dots, l_1\}$, $\{l_1 + 1, \dots, 2l_1\}$, \dots , $\{(e_1 - 1)l_1 + 1, \dots, e_1 l_1\}$.

It is straightforward but tedious to check that $H \cap K^{g^{-1}}$ coincides with the subgroup of $\Sigma_{e_1 l_1}$ given above. \square

Lemma 4.10. Let $\Gamma_1, \dots, \Gamma_s$ be finite groups, $d_i \in B(\Gamma_i)$ for $i = 1, \dots, s$ and $e \in \mathbb{N}$. Let $\bar{\phi}$ be the group homomorphism

$$\Sigma_e \langle \Gamma_1 \times \cdots \times \Gamma_s \rangle \rightarrow \Sigma_e \langle \Gamma_1 \rangle \times \cdots \times \Sigma_e \langle \Gamma_s \rangle.$$

Then, $\bar{\phi}^*(\bigotimes_{i=1}^s T_e(d_i)) = T_e(\bigotimes_{i=1}^s d_i)$.

Proof. First we establish some notations:


$$G = \Sigma_e \langle \Gamma_1 \rangle \times \cdots \times \Sigma_e \langle \Gamma_s \rangle.$$

$$K_j = \Sigma_j \langle \Gamma_2 \rangle \times \cdots \times \Gamma_s, \quad j = 0, 1, \dots, e.$$

$$\bar{K}_j = \Sigma_j \langle \Gamma_1 \rangle \times \cdots \times \Gamma_s, \quad j = 0, 1, \dots, e.$$

$$H_j = (\Sigma_j \langle \Gamma_1 \rangle \times \Sigma_{e-j} \langle \Gamma_1 \rangle) \times \Sigma_e \langle \Gamma_2 \rangle \times \cdots \times \Sigma_e \langle \Gamma_s \rangle, \quad j = 0, 1, \dots, s.$$

In the following diagram all homomorphisms behave as expected:



Observe that G has only one $(\psi(H_j), \bar{\phi}(\bar{K}_e))$ -double coset.

Now, applying the Mackey theorem and using the commutativity of the diagram and Proposition (3.3)(iv), one has

$$\bar{\phi}^* \psi_* = \beta_* \gamma_* \delta^* \tau^* \alpha^* \phi^* \quad (3)$$

Now, let us prove the lemma.

If $d_1 = X_i$ is a Γ_i -set for all $i = 1, \dots, s$, then it is easy to check the result. We will prove it for $d_1 = X_1 - X'_1$, $d_2 = X_2, \dots, d_s = X_s$ with X_i a Γ_i -set ($i = 1, \dots, s$) and X'_1 a Γ_1 -set. With a similar process we would do it after that for any $d_2 \in B(\Gamma_2)$, and so on.

Put $z := \bar{\phi}^*[T_e(d_1) \otimes T_e(X_2) \otimes \cdots \otimes T_e(X_s)]$.

Now,

$$\begin{aligned} & \bar{\phi}^*[T_e(X_1) \otimes T_e(X_2) \otimes \cdots \otimes T_e(X_s)] \\ &= \sum_{j=0}^e \bar{\phi}^*[(T_j(d_1) \times T_{e-j}(X'_1)) \otimes T_e(X_2) \otimes \cdots \otimes T_e(X_s)] \end{aligned}$$

$$= z + \sum_{j=0}^{e-1} \bar{\phi}^* \psi_* [(T_j(d_1) \otimes T_{e-j}(X'_1)) \otimes T_e(X_2) \otimes \cdots \otimes T_e(X_s)]$$

by (3) and the inductive hypothesis on s

$$= z + \sum_{j=0}^{e-1} \beta_* \gamma_* \delta^* \tau^* \alpha^* [(T_j(d_1) \otimes T_{e-j}(X'_1)) \otimes T_e(X_2 \otimes \cdots \otimes X_s)]$$

by Proposition 4.2(iii)

$$\begin{aligned} &= z + \sum_{j=0}^{e-1} \beta_* \gamma_* \delta^* \tau^* [(T_j(d_1) \otimes T_{e-j}(X'_1)) \otimes (T_j(X_2 \otimes \cdots \otimes X_s) \\ &\quad \otimes T_{e-j}(X_2 \otimes \cdots \otimes X_s))] \\ &= z + \sum_{j=0}^{e-1} \beta_* \gamma_* \delta^* [(T_j(d_1) \otimes T_j(X_2 \otimes \cdots \otimes X_s)) \otimes (T_{e-j}(X'_1) \\ &\quad \otimes T_{e-j}(X_2 \otimes \cdots \otimes X_s))] \end{aligned}$$

by the inductive hypothesis on e for $s=2$

$$= z + \sum_{j=0}^{e-1} \beta_* \gamma_* [T_j(d_1 \otimes X_2 \otimes \cdots \otimes X_s) \otimes T_{e-j}(X'_1 \otimes X_2 \otimes \cdots \otimes X_s)]$$

by the definition of the cross product

$$= z + \sum_{j=0}^{e-1} [T_j(d_1 \times X_2 \times \cdots \times X_s) \times T_{e-j}(X'_1 \times X_2 \times \cdots \times X_s)].$$

On the other hand,

$$\begin{aligned} &\bar{\phi}^* (T_e(X_1) \otimes \cdots \otimes T_e(X_s)) = T_e(X_1 \otimes \cdots \otimes X_s) \\ &= \sum_{j=0}^e [T_j(d_1 \otimes X_2 \otimes \cdots \otimes X_s) \times T_{e-j}(X'_1 \otimes X_2 \otimes \cdots \otimes X_s)]. \end{aligned}$$

Thus,

$$z = T_e(d_1 \otimes X_2 \otimes \cdots \otimes X_s). \quad \square$$

Notations for the remainder of Section 4 (except Lemmas 4.15 and 4.18)

$\varepsilon = (\varepsilon_1, \dots, \varepsilon_r)$ is an r -partition of k .

$k(\varepsilon) := \varepsilon_1 l_1 + \cdots + \varepsilon_r l_r$.

$\omega(\varepsilon) = (\omega(\varepsilon)_1, \dots, \omega(\varepsilon)_s)$ is an s -partition of $k(\varepsilon)$.

$\theta'_\varepsilon : \Sigma_{\varepsilon_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{\varepsilon_r} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{\varepsilon_1 l_1} \times \cdots \times \Sigma_{\varepsilon_r l_r}$.

$\theta''_\varepsilon : \Sigma_{\varepsilon_1 l_1} \times \cdots \times \Sigma_{\varepsilon_r l_r} \rightarrow \Sigma_{k(\varepsilon)}$.

$\theta_\varepsilon = \theta''_\varepsilon \theta'_\varepsilon$.

$\pi_\varepsilon : \Sigma_{\varepsilon_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{\varepsilon_r} \langle \Sigma_{l_r} \rangle \rightarrow \Sigma_{\varepsilon_1} \times \cdots \times \Sigma_{\varepsilon_r}$.

$\delta_\varepsilon : \Sigma_{\varepsilon_1} \times \cdots \times \Sigma_{\varepsilon_r} \rightarrow \Sigma_k$.

$L(\varepsilon, b, a) = (\bigotimes_{j=1}^r T_{\varepsilon_j}(a_j))(\pi_\varepsilon^* \delta_\varepsilon^*(b))$.

$\theta'_{\omega(\varepsilon)} : \Sigma_{\omega(\varepsilon)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\omega(\varepsilon)_s} \langle \Sigma_{m_s} \rangle \rightarrow \Sigma_{\omega(\varepsilon)_1 m_1} \times \cdots \times \Sigma_{\omega(\varepsilon)_s m_s}$.

$\theta''_{\omega(\varepsilon)} : \Sigma_{\omega(\varepsilon)_1 m_1} \times \cdots \times \Sigma_{\omega(\varepsilon)_s m_s} \rightarrow \Sigma_{\omega(\varepsilon)_1 m_1 + \cdots + \omega(\varepsilon)_s m_s}$.

$$\theta_{\omega(\varepsilon)} = \theta''_{\omega(\varepsilon)} \theta'_{\omega(\varepsilon)}.$$

$$\pi_{\omega(\varepsilon)} : \Sigma_{\omega(\varepsilon)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\omega(\varepsilon)_s} \langle \Sigma_{m_s} \rangle \rightarrow \Sigma_{\omega(\varepsilon)_1} \times \cdots \times \Sigma_{\omega(\varepsilon)_s}.$$

$$\delta_{\omega(\varepsilon)} : \Sigma_{\omega(\varepsilon)_1} \times \cdots \times \Sigma_{\omega(\varepsilon)_s} \rightarrow \Sigma_{k(\varepsilon)}.$$

$\{\gamma(j)^1, \dots, \gamma(j)^{p_j}\}$ is the set of all s -partitions of l_j .

$$\gamma(j)^i = (\gamma(j)_1^i, \dots, \gamma(j)_s^i).$$

$$t := p_1 + \cdots + p_r.$$

$A = (a_{ij})$ is an \mathbb{N} -matrix $r \times s$ with row sums $(\varepsilon_1 l_1, \dots, \varepsilon_r l_r)$ and column sums $(\omega(\varepsilon)_1, \dots, \omega(\varepsilon)_s)$.

B_{ε_j} is an \mathbb{N} -matrix $\varepsilon_j \times s$ with row sums (l_j, \dots, l_j) and column sums (a_{j1}, \dots, a_{js}) .

The classes $[B_{\varepsilon_j}]$ are defined as in Lemma 4.9.

B is the ‘diagonal’ sum of $[B_{\varepsilon_j}]$ ($j=1, \dots, r$).

$$B = \left[\begin{array}{ccc} [B_{\varepsilon_1}] & & 0 \\ & [B_{\varepsilon_2}] & \\ 0 & & \ddots \\ & & & \ddots \end{array} \right] \left. \begin{array}{l} \} \varepsilon_1 \\ \} \varepsilon_2 \\ \vdots \end{array} \right\}$$

$\underbrace{\quad}_S \quad \underbrace{\quad}_S \quad \dots$

$$\mu_A : (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}})$$

$$\rightarrow (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{1s}}) \times \cdots \times (\Sigma_{a_{r1}} \times \cdots \times \Sigma_{a_{rs}}).$$

$$\varrho_A : (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{1s}}) \times \cdots \times (\Sigma_{a_{r1}} \times \cdots \times \Sigma_{a_{rs}}) \rightarrow \Sigma_{\varepsilon_1 l_1} \times \cdots \times \Sigma_{\varepsilon_r l_r}.$$

$$\phi_A : (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}}) \rightarrow \Sigma_{\omega(\varepsilon)_1} \times \cdots \times \Sigma_{\omega(\varepsilon)_s}.$$

$e_{p_1} + \cdots + e_{p_{j-1}+q}$ ($1 \leq q \leq p_j$) is the number of rows of B_{ε_j} which are equal to $\gamma(j)^q$.

$$H_B = (\Sigma_{e_1} \langle \Sigma_{\gamma(1)_1^1} \rangle \times \cdots \times \Sigma_{\gamma(1)_s^1} \rangle) \times \cdots \times \Sigma_{e_{p_1}} \langle \Sigma_{\gamma(1)_1^{p_1}} \rangle \times \cdots \times \Sigma_{\gamma(1)_s^{p_1}} \rangle) \times \cdots$$

$$\times \cdots \times (\Sigma_{e_{t-p_r+1}} \langle \Sigma_{\gamma(r)_1^1} \rangle \times \cdots \times \Sigma_{\gamma(r)_s^1} \rangle) \times \cdots \times \Sigma_{e_t} \langle \Sigma_{\gamma(r)_1^{p_r}} \rangle \times \cdots \times \Sigma_{\gamma(r)_s^{p_r}} \rangle).$$

$H_B \langle \Sigma_{m_i} \rangle$: put $\Sigma_{\gamma(j)_i^q} \langle \Sigma_{m_i} \rangle$ instead of $\Sigma_{\gamma(j)_i^q}$ in H_B for $j=1, \dots, r$; $i=1, \dots, s$; $q=1, \dots, p_j$.

$$\eta_B : H_B \rightarrow \Sigma_{\varepsilon_1} \langle \Sigma_{l_1} \rangle \times \cdots \times \Sigma_{\varepsilon_r} \langle \Sigma_{l_r} \rangle.$$

$\tau : H_B^g \rightarrow H_B$ is conjugation by g^{-1} , where $g = g_1 \times \cdots \times g_r$ and g_j is a representative of $\Psi([B_{\varepsilon_j}])$ in Lemma 4.9 and defined in the same way.

$$\nu_B : H_B^g \hookrightarrow (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{1s}}) \times \cdots \times (\Sigma_{a_{r1}} \times \cdots \times \Sigma_{a_{rs}}).$$

$H_B^g \langle \Sigma_{m_i} \rangle$ is the subgroup of

$$\begin{aligned} & [(\Sigma_{e_1} \langle \Sigma_{\gamma(1)_1^1} \rangle \times \cdots \times \Sigma_{e_{p_1}} \langle \Sigma_{\gamma(1)_1^{p_1}} \rangle) \langle \Sigma_{m_1} \rangle \times \cdots \\ & \times (\Sigma_{e_1} \langle \Sigma_{\gamma(1)_s^1} \rangle \times \cdots \times \Sigma_{e_{p_1}} \langle \Sigma_{\gamma(1)_s^{p_1}} \rangle) \langle \Sigma_{m_s} \rangle] \times \cdots \\ & \times [(\Sigma_{e_{t-p_r+1}} \langle \Sigma_{\gamma(r)_1^1} \rangle \times \cdots \times \Sigma_{e_t} \langle \Sigma_{\gamma(r)_1^{p_r}} \rangle) \langle \Sigma_{m_1} \rangle \times \cdots \\ & \times (\Sigma_{e_{t-p_r+1}} \langle \Sigma_{\gamma(r)_s^1} \rangle \times \cdots \times \Sigma_{e_t} \langle \Sigma_{\gamma(r)_s^{p_r}} \rangle) \langle \Sigma_{m_s} \rangle] \end{aligned}$$

consisting of all elements such that the ‘component’ in Σ_{e_i} is the same for all places where it occurs ($i=1, \dots, t$).

$$\tilde{\varphi}_A : (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \langle \Sigma_{m_1} \rangle \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}}) \langle \Sigma_{m_s} \rangle$$

$$\rightarrow \Sigma_{\omega(\varepsilon)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\omega(\varepsilon)_s} \langle \Sigma_{m_s} \rangle.$$

$$\tilde{\pi}_{\omega(\varepsilon)} : (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \langle \Sigma_{m_1} \rangle \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}}) \langle \Sigma_{m_s} \rangle$$

$$\rightarrow (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}}).$$

$$\begin{aligned}
\bar{v}_B &: H_B^g \langle \Sigma_{m_i} \rangle \hookrightarrow (\Sigma_{a_{11}} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{a_{1s}} \langle \Sigma_{m_s} \rangle) \times \cdots \times (\Sigma_{a_{r1}} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{a_{rs}} \langle \Sigma_{m_s} \rangle). \\
\bar{\mu}_A &: (\Sigma_{a_{11}} \times \cdots \times \Sigma_{a_{r1}}) \langle \Sigma_{m_1} \rangle \times \cdots \times (\Sigma_{a_{1s}} \times \cdots \times \Sigma_{a_{rs}}) \langle \Sigma_{m_s} \rangle \\
&\rightarrow (\Sigma_{a_{11}} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{a_{1s}} \langle \Sigma_{m_s} \rangle) \times \cdots \times (\Sigma_{a_{r1}} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{a_{rs}} \langle \Sigma_{m_s} \rangle). \\
\pi_B &: H_B \langle \Sigma_{m_i} \rangle \rightarrow H_B. \\
\bar{\pi}_B &: H_B^g \langle \Sigma_{m_i} \rangle \rightarrow H_B^g. \\
\sigma &= (\sigma_1, \dots, \sigma_t) \text{ is a } t\text{-partition of } k. \\
\gamma(h) &= \gamma(j)^q \text{ if } h = p_1 + \cdots + p_{j-1} + q \text{ with } 1 \leq q \leq p_j. \\
n_h &= \gamma(h)_1 m_1 + \cdots + \gamma(h)_s m_s. \\
\theta'_{\gamma(h)} &: \Sigma_{\gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\gamma(h)_s} \langle \Sigma_{m_s} \rangle \rightarrow \Sigma_{\gamma(h)_1 m_1} \times \cdots \times \Sigma_{\gamma(h)_s m_s}. \\
\theta''_{\gamma(h)} &: \Sigma_{\gamma(h)_1 m_1} \times \cdots \times \Sigma_{\gamma(h)_s m_s} \rightarrow \Sigma_{n_h}. \\
\theta_{\gamma(h)} &= \theta''_{\gamma(h)} \theta'_{\gamma(h)}. \\
\delta_{\gamma(h)} &: \Sigma_{\gamma(h)_1} \times \cdots \times \Sigma_{\gamma(h)_s} \rightarrow \Sigma_{l_j} \text{ if } \gamma(h) = \gamma(j)^q. \\
\pi_{\gamma(h)} &: \Sigma_{\gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\gamma(h)_s} \langle \Sigma_{m_s} \rangle \rightarrow \Sigma_{\gamma(h)_1} \times \cdots \times \Sigma_{\gamma(h)_s}. \\
L(\gamma(h), a_j, c) &= (\bigotimes_{i=1}^s T_{\gamma(h)_i}(c_i)) (\pi_{\gamma(h)}^* \delta_{\gamma(h)}^*(a_j)). \\
\theta'_\sigma &: \Sigma_{\sigma_1} \langle \Sigma_{n_1} \rangle \times \cdots \times \Sigma_{\sigma_t} \langle \Sigma_{n_t} \rangle \rightarrow \Sigma_{\sigma_1 n_1} \times \cdots \times \Sigma_{\sigma_t n_t}. \\
\theta''_\sigma &: \Sigma_{\sigma_1 n_1} \times \cdots \times \Sigma_{\sigma_t n_t} \rightarrow \Sigma_{\sigma_1 n_1 + \cdots + \sigma_t n_t}. \\
\theta_\sigma &= \theta''_\sigma \theta'_\sigma. \\
\pi_\sigma &: \Sigma_{\sigma_1} \langle \Sigma_{n_1} \rangle \times \cdots \times \Sigma_{\sigma_t} \langle \Sigma_{n_t} \rangle \rightarrow \Sigma_{\sigma_1} \times \cdots \times \Sigma_{\sigma_t}. \\
\delta_\sigma &: \Sigma_{\sigma_1} \times \cdots \times \Sigma_{\sigma_t} \rightarrow \Sigma_k. \\
\bar{\theta}_{\gamma(h)} &: \Sigma_{\sigma_h} \langle (\Sigma_{\gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\gamma(h)_s} \langle \Sigma_{m_s} \rangle) \rangle \rightarrow \Sigma_{\sigma_h} \langle \Sigma_{n_h} \rangle. \\
\bar{\delta}_{\gamma(h)} &: \Sigma_{\sigma_h} \langle \Sigma_{\gamma(h)_1} \times \cdots \times \Sigma_{\gamma(h)_s} \rangle \rightarrow \Sigma_{\sigma_h} \langle \Sigma_{l_j} \rangle. \\
\bar{\pi}_{\gamma(h)} &: \Sigma_{\sigma_h} \langle (\Sigma_{\gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\gamma(h)_s} \langle \Sigma_{m_s} \rangle) \rangle \rightarrow \Sigma_{\sigma_h} \langle \Sigma_{\gamma(h)_1} \times \cdots \times \Sigma_{\gamma(h)_s} \rangle. \\
\bar{\tau} &: H_B^g \langle \Sigma_{m_i} \rangle \rightarrow H_B \langle \Sigma_{m_i} \rangle \text{ defined from } \tau \text{ in the obvious way.} \\
\psi_h^1 &: \Sigma_{\sigma_h} \langle (\Sigma_{\gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\gamma(h)_s} \langle \Sigma_{m_s} \rangle) \rangle \rightarrow \Sigma_{\sigma_h} \langle \Sigma_{\gamma(h)_1} \langle \Sigma_{m_1} \rangle \rangle \times \cdots \times \Sigma_{\sigma_h} \langle \Sigma_{\gamma(h)_s} \langle \Sigma_{m_s} \rangle \rangle. \\
\psi_h^2 &: (\text{codomain}(\psi_h^1)) \rightarrow \Sigma_{\sigma_h \gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\sigma_h \gamma(h)_s} \langle \Sigma_{m_s} \rangle. \\
\psi^3 &: \times_{h=1}^t [\Sigma_{\sigma_h \gamma(h)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\sigma_h \gamma(h)_s} \langle \Sigma_{m_s} \rangle] \\
&\rightarrow [\Sigma_{\sigma_1 \gamma(1)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\sigma_1 \gamma(1)_s} \langle \Sigma_{m_s} \rangle] \times \cdots \times [\Sigma_{\sigma_t \gamma(t)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\sigma_t \gamma(t)_s} \langle \Sigma_{m_s} \rangle]. \\
\psi^4 &: (\text{codomain}(\psi^3)) \rightarrow \Sigma_{\omega(\varepsilon)_1} \langle \Sigma_{m_1} \rangle \times \cdots \times \Sigma_{\omega(\varepsilon)_s} \langle \Sigma_{m_s} \rangle. \\
\eta'_j &= \bigotimes_{h_j} \bar{\delta}_{\gamma(h_j)} \text{ where } h_j = p_1 + \cdots + p_{j-1} + 1, \dots, p_1 + \cdots + p_j \text{ for } j = 1, \dots, r. \\
\eta''_j &: \Sigma_{\sigma_{p_1} + \cdots + p_{j-1} + 1} \langle \Sigma_{l_j} \rangle \times \cdots \times \Sigma_{\sigma_{p_1 + \cdots + p_j}} \langle \Sigma_{l_j} \rangle \rightarrow \Sigma_{\varepsilon_j} \langle \Sigma_{l_j} \rangle.
\end{aligned}$$

Proposition 4.11. *Property 3 of Definition 4.4 is valid in $B(S)$.*

Proof. Let $b \in B(\Sigma_k)$, $a = a_1 + \cdots + a_r$ with $a_j \in B(\Sigma_{l_j})$ ($j = 1, \dots, r$), $c = c_1 + \cdots + c_s$ with $c_i \in B(\Sigma_{m_i})$ ($i = 1, \dots, s$).

We want to prove $(b \vee a) \vee c = b \vee (a \vee c)$.

$$\begin{aligned}
(b \vee a) \vee c &= \left[\sum_{\varepsilon} (\theta_{\varepsilon})_* L(\varepsilon, b, a) \right] \vee c \\
&= \sum_{\varepsilon} \sum_{\omega(\varepsilon)} (\theta_{\omega(\varepsilon)})_* \left[\left(\bigotimes_{i=1}^s T_{\omega(\varepsilon)_i}(c_i) \right) (\pi_{\omega(\varepsilon)}^* \delta_{\omega(\varepsilon)}^* (\theta_{\varepsilon})_* L(\varepsilon, b, a)) \right]
\end{aligned}$$

by the Mackey theorem and Lemma 4.7 (put $U = (\bigotimes_{i=1}^s T_{\omega(\varepsilon)_i}(c_i))$)

$$= \sum_{\varepsilon} \sum_{\omega(\varepsilon)} (\theta_{\omega(\varepsilon)})_* \left[U \left(\pi_{\omega(\varepsilon)}^* \left[\sum_A (\phi_A)_* \mu_A^* \varrho_A^* (\theta_{\varepsilon}')_* L(\varepsilon, b, a) \right] \right) \right]$$

by the Mackey theorem and Lemma 4.9

$$= \sum_{\varepsilon} \sum_{\omega(\varepsilon)} \sum_A (\theta_{\omega(\varepsilon)})_* \left[U \left(\pi_{\omega(\varepsilon)}^* (\phi_A)_* \mu_A^* \left[\sum_B (v_B)_* \tau^* \eta_B^* L(\varepsilon, b, a) \right] \right) \right] =$$

by Proposition 3.6: $\pi_{\omega(\varepsilon)}^* (\phi_A)_* = (\bar{\phi}_A)_* \bar{\pi}_{\omega(\varepsilon)}^*$; so, $\bar{\pi}_{\omega(\varepsilon)}^* \mu_A^* (v_B)_* = \bar{\pi}_{\omega(\varepsilon)}^* (\mu_A^{-1})_* (v_B)_* = \bar{\pi}_{\omega(\varepsilon)}^* (\mu_A^{-1} v_B)_* = (\bar{\mu}_A^{-1})_* (\bar{v}_B)_* \bar{\pi}_B^*$

$$= \sum_{\varepsilon} \sum_{\omega(\varepsilon)} \sum_A \sum_B (\theta_{\omega(\varepsilon)})_* [U((\bar{\phi}_A)_* (\bar{\mu}_A^{-1})_* (\bar{v}_B)_* \bar{\pi}_B^* \tau^* \eta_B^* L(\varepsilon, b, a))]$$

by Lemma 4.12 and Frobenius reciprocity

$$= \sum_{\sigma \in D} (\theta_{\omega(\varepsilon)} \bar{\phi}_A \bar{\mu}_A^{-1} \bar{v}_B)_* [[\bar{v}_B^* (\bar{\mu}_A^{-1})^* \bar{\phi}_A^* (U)] [\bar{\pi}_B^* \tau^* \eta_B^* L(\varepsilon, b, a)]]$$

by Lemma 4.13 and $\zeta_* = \text{id}$

$$= \sum_{\sigma} \left(\theta_{\sigma} \left(\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)} \right) \bar{\tau} \right)_* [[\bar{v}_B^* (\bar{\mu}_A^{-1})^* \bar{\phi}_A^* (U)] [\bar{\pi}_B^* \tau^* \eta_B^* L(\varepsilon, b, a)]]$$

by Lemma 4.14

$$= \sum_{\sigma} \left(\theta_{\sigma} \left(\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)} \right) \bar{\tau} \right)_* \left[\left(\bar{\tau}^* \left[\bigotimes_{h=1}^l T_{\sigma_h} \left(\bigotimes_{i=1}^s T_{\gamma(h)_i} (c_i) \right) \right] \right) \right. \\ \left. (\bar{\pi}_B^* \tau^* \eta_B^* L(\varepsilon, b, a)) \right] =$$

by Proposition 3.3 (put $\bigotimes_{h=1}^l T_{\sigma_h} (\bigotimes_{i=1}^s T_{\gamma(h)_i} (c_i)) = V$ and $\theta_{\sigma} (\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)} \bar{\tau} = \Lambda)$)

$$= \sum_{\sigma} \Lambda_* \left[\bar{\tau}^* (V) \left(\bar{\pi}_B^* \tau^* \eta_B^* \left[\bigotimes_{j=1}^r T_{\varepsilon_j} (a_j) \right] \right) (\bar{\pi}_B^* \tau^* \eta_B^* \pi_{\varepsilon}^* \delta^* (b)) \right]$$

by Lemma 4.16

$$= \sum_{\sigma} \Lambda_* \left[\bar{\tau}^* (V) \left(\bar{\tau}^* \left[\bigotimes_{h=1}^l T_{\sigma_h} (\pi_{\gamma(h)}^* \delta_{\gamma(h)}^* (a_h)) \right] \right) (\bar{\pi}_B^* \tau^* \eta_B^* \pi_{\varepsilon}^* \delta_{\varepsilon}^* (b)) \right]$$

by Lemma 4.17 and Proposition 3.3(i), (iv) applied to $\bar{\tau}$

$$= \sum_{\sigma} (\theta_{\sigma})_* \left(\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)} \right)_* \left[V \left(\bigotimes_{h=1}^l T_{\sigma_h} (\pi_{\gamma(h)}^* \delta_{\gamma(h)}^* (a_h)) \right) \right. \\ \left. \left(\left(\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)} \right)^* \pi_{\sigma}^* \delta_{\sigma}^* (b) \right) \right]$$

by Proposition 4.2(iv) (remember $V = \dots$) and Frobenius reciprocity

$$\begin{aligned}
&= \sum_{\sigma} (\theta_{\sigma})_* \left[\left(\bigotimes_{h=1}^t (\bar{\theta}_{\gamma(h)})_* T_{\sigma_h} [L(\gamma(h), a_h, c)] \right) (\pi_{\sigma}^* \delta_{\sigma}^*(b)) \right] \\
&\quad \text{by Lemma 4.18} \\
&= \sum_{\sigma} (\theta_{\sigma})_* \left[\left(\bigotimes_{h=1}^t T_{\sigma_h} [(\theta_{\gamma(h)})_* L(\gamma(h), a_h, c)] \right) (\pi_{\sigma}^* \delta_{\sigma}^*(b)) \right] \\
&= b \vee (a \vee c).
\end{aligned}$$

The last equality is due to

$$\begin{aligned}
a \vee c &= \sum_{j=1}^r a_j \vee c = \sum_{j=1}^r \left[\sum_{h_j} (\theta_{\gamma(h_j)})_* L(\gamma(h_j), a_j, c) \right] \\
&= \sum_{h=1}^t (\theta_{\gamma(h)})_* L(\gamma(h), a_h, c)
\end{aligned}$$

where $h_j = p_1 + \dots + p_{j-1} + 1, \dots, p_1 + \dots + p_j$ and $a_h = a_j$ when h is equal to some h_j . \square

Lemma 4.12. Let $D = \{\sigma = (\sigma_1, \dots, \sigma_t): \sigma \text{ a } t\text{-partition of } k\}$ and $M = \{B: B \text{ as defined in notations}\}$.

Then there exists a bijective map $\chi: D \rightarrow M$.

Proof. For $\sigma \in D$ we define $\chi(\sigma) = B$, the ‘diagonal sum’ of $[B_{\varepsilon_j}]$ ($j = 1, \dots, r$) with B_{ε_j} the matrix consisting of σ_h rows equal to $\gamma(j)^q$, where $h = p_1 + \dots + p_{j-1} + q$ and $1 \leq q \leq p_j$.

Mapping each matrix B to the partition σ which has $\sigma_h =$ number of rows of the j th ‘diagonal’ component of B , (B_{ε_j}) , equal to $\gamma(j)^q$ ($h = p_1 + \dots + p_{j-1} + q$) we obtain an inverse map of χ . \square


Observations.

$$\text{card } D = \text{card } M = \sum_{\varepsilon} \begin{bmatrix} p_1 \\ \varepsilon_1 \end{bmatrix} \dots \begin{bmatrix} p_r \\ \varepsilon_r \end{bmatrix},$$

where $\varepsilon = (\varepsilon_1, \dots, \varepsilon_r)$ ranges over all r -partitions of k and $\binom{p_j}{\varepsilon_j}$ is the number of combinations of p_j elements, ε_j at a time, when repetitions are allowed.

Note that now σ determines an r -partition of k : $\varepsilon = (\varepsilon_1, \dots, \varepsilon_r)$ with $\varepsilon_j = \sigma_{p_1 + \dots + p_{j-1} + 1} + \dots + \sigma_{p_1 + \dots + p_j}$, a matrix A whose rows are the column sums of the matrices B_{ε_j} , an s -partition of $k(\varepsilon)$: $\omega(\varepsilon)$ given by the column sums of A , and obviously a matrix B .

The following picture may be useful:



Lemma 4.13. *There exists an inner automorphism ζ of the group $\Sigma_{\omega(\varepsilon)_1 m_1 + \dots + \omega(\varepsilon)_s m_s}$ such that*

$$\zeta \theta_\sigma \left(\bigotimes_{h=1}^t \bar{\theta}_{\gamma(h)} \right) \bar{\tau} = \theta_{\omega(\varepsilon)} \bar{\phi}_A \bar{\mu}_A^{-1} \bar{v}_B.$$

Proof. First, observe that

$$\sigma_1 n_1 + \dots + \sigma_t n_t = \omega(\varepsilon)_1 m_1 + \dots + \omega(\varepsilon)_s m_s.$$

Now, forgetting ζ , both compositions are homomorphisms from $H_B^g \langle \Sigma_{m_i} \rangle$ to $\Sigma_{\sigma_1 n_1 + \dots + \sigma_t n_t}$ and all components are either inclusions or conjugations in $\Sigma_{\sigma_1 n_1 + \dots + \sigma_t n_t}$. \square

Lemma 4.14.

$$\bar{v}_B^* (\bar{\mu}_A^{-1})^* \bar{\phi}_A^* \left(\bigotimes_{i=1}^s T_{\omega(\varepsilon)_i} (c_i) \right) = \bar{\tau}^* \left[\bigotimes_{h=1}^t T_{\sigma_h} \left(\bigotimes_{i=1}^s T_{\gamma(h)_i} (c_i) \right) \right].$$

Proof. It is not difficult to check that

$$\bar{\phi}_A \bar{\mu}_A^{-1} \bar{v}_B = \psi^4 \psi^3 \left(\bigotimes_{h=1}^t \psi_h^2 \right) \left(\bigotimes_{h=1}^t \psi_h^1 \right) \bar{\tau}.$$

Now,

$$\begin{aligned} & \bar{v}_B^* (\bar{\mu}_A^{-1})^* \bar{\phi}_A^* \left(\bigotimes_{i=1}^s T_{\omega(\varepsilon)_i} (c_i) \right) \\ &= \bar{\tau}^* \left(\bigotimes_{h=1}^t \psi_h^1 \right)^* \left(\bigotimes_{h=1}^t \psi_h^2 \right)^* (\psi^3)^* (\psi^4)^* \left[\bigotimes_{i=1}^s T_{\omega(\varepsilon)_i} (c_i) \right] \end{aligned}$$

by Proposition 4.2(iii)

$$\begin{aligned} &= \bar{\tau}^* \left(\bigotimes_{h=1}^l \psi_h^1 \right)^* \left(\bigotimes_{h=1}^l \psi_h^2 \right)^* (\psi^3)^* \left[\bigotimes_{i=1}^s \left(\bigotimes_{h=1}^l T_{\sigma_h \gamma(h)_i}(c_i) \right) \right] \\ &= \bar{\tau}^* \left(\bigotimes_{h=1}^l \psi_h^1 \right)^* \left(\bigotimes_{h=1}^l \psi_h^2 \right)^* \left[\bigotimes_{h=1}^l \left(\bigotimes_{i=1}^s T_{\sigma_h \gamma(h)_i}(c_i) \right) \right] \end{aligned}$$

by Proposition 4.2(v)

$$= \bar{\tau}^* \left[\bigotimes_{h=1}^l (\psi_h^1)^* \left(\bigotimes_{i=1}^s T_{\sigma_h} T_{\gamma(h)_i}(c_i) \right) \right]$$

by Lemma 4.10

$$= \bar{\tau}^* \left[\bigotimes_{h=1}^l T_{\sigma_h} \left(\bigotimes_{i=1}^s T_{\gamma(h)_i}(c_i) \right) \right]. \quad \square$$

Lemma 4.15. Let $\Psi: \Gamma \rightarrow \Omega$ be a group homomorphism and $\bar{\Psi}_k = \Sigma_k \langle \Psi \rangle: \Sigma_k \langle \Gamma \rangle \rightarrow \Sigma_k \langle \Omega \rangle$ ($k \in \mathbb{N}$). Then

$$T_k \Psi^* = \bar{\Psi}^* T_k.$$

Proof. The following diagram commutes for Ω -sets:

$$\begin{array}{ccc} B(\Omega) & \xrightarrow{\Psi^*} & B(\Gamma) \\ T \downarrow & & \downarrow T \\ \prod_{k \geq 0} B(\Sigma_k \langle \Omega \rangle) & \xrightarrow{\prod_{k \geq 0} \bar{\Psi}_k^*} & \prod_{k \geq 0} B(\Sigma_k \langle \Gamma \rangle) \end{array}$$

$T\Psi^*$ is a homomorphism from ‘+’ to ‘ \times ’. Finally, $(\prod_{k \geq 0} \Psi_k^*)T$ is a homomorphism from ‘+’ to ‘ \times ’ because $\prod_{k \geq 0} \Psi_k^*$ is from ‘ \times ’ to ‘ \times ’ since the following diagram commutes:

$$\begin{array}{ccc} B(\Sigma_i \langle \Omega \rangle) & \xrightarrow{\bar{\Psi}_i^*} & B(\Sigma_i \langle \Gamma \rangle) \\ \uparrow \text{‘}\times\text{’} & & \uparrow \text{‘}\times\text{’} \\ B(\Sigma_i \langle \Omega \rangle) \otimes B(\Sigma_j \langle \Omega \rangle) & \xrightarrow{\Psi_i^* \otimes \Psi_j^*} & B(\Sigma_i \langle \Gamma \rangle) \otimes B(\Sigma_j \langle \Gamma \rangle) \end{array}$$

Lemma 4.16.

$$\pi_B^* \tau^* \eta_B^* \left(\bigotimes_{j=1}^r T_{e_j}(a_j) \right) = \bar{\tau}^* \left[\bigotimes_{h=1}^l T_{\sigma_h} (\pi_{\gamma(h)}^* \delta_{\gamma(h)}^*(a_h)) \right]$$

where $a_h = a_j$ if $h \in \{p_1 + \dots + p_{j-1} + 1, \dots, p_1 + \dots + p_j\}$.

Proof. $\eta_B = (\bigotimes_{j=1}^r \eta_j'')(\bigotimes_{j=1}^r \eta_j')$. So,

$$\begin{aligned}
 \bar{\pi}_B^* \tau^* \eta_B^* \left(\bigotimes_{j=1}^r T_{\varepsilon_j}(a_j) \right) &= \bar{\pi}_B^* \tau^* \left(\bigotimes_{j=1}^r \eta_j' \right)^* \left[\bigotimes_{j=1}^r (\eta_j'')^* T_{\varepsilon_j}(a_j) \right] \\
 &\quad \text{by Proposition 4.2(iii) and putting } h_j = p_1 + \dots + p_{j-1} + 1, \dots, \\
 &\quad p_1 + \dots + p_j \\
 &= \bar{\pi}_B^* \tau^* \left(\bigotimes_{j=1}^r \eta_j' \right)^* \left[\bigotimes_{j=1}^r \left(\bigotimes_{h_j} T_{\sigma_{h_j}}(a_j) \right) \right] \\
 &= \bar{\pi}_B^* \tau^* \left[\bigotimes_{j=1}^r \left(\bigotimes_{h_j} \delta_{\gamma(h_j)}^* T_{\sigma_{h_j}}(a_j) \right) \right] = \bar{\pi}_B^* \tau^* \left[\bigotimes_{h=1}^l \delta_{\gamma(h)}^* T_{\sigma_h}(a_h) \right] \\
 &\quad \text{by } \tau \bar{\pi}_B = \pi_B \bar{\tau} \\
 &= \bar{\tau}^* \pi_B^* \left[\bigotimes_{h=1}^l \delta_{\gamma(h)}^* T_{\sigma_h}(a_h) \right] = \bar{\tau}^* \left[\bigotimes_{h=1}^l \pi_{\gamma(h)}^* \delta_{\gamma(h)}^* T_{\sigma_h}(a_h) \right] \\
 &\quad \text{by Lemma 4.15} \\
 &= \bar{\tau}^* \left[\bigotimes_{h=1}^l T_{\sigma_h}(\pi_{\gamma(h)}^* \delta_{\gamma(h)}^*(a_h)) \right]. \quad \square
 \end{aligned}$$

Lemma 4.17. $\delta_\varepsilon \pi_\varepsilon \eta_B \tau \bar{\pi}_B = \delta_\sigma \pi_\sigma (\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)}) \bar{\tau}$.

Proof. $\tau \bar{\pi}_B = \pi_B \bar{\tau}$, and then it is easy to check that

$$\delta_\varepsilon \pi_\varepsilon \eta_B \pi_B = \delta_\sigma \pi_\sigma \left(\bigotimes_{h=1}^l \bar{\theta}_{\gamma(h)} \right). \quad \square$$

Lemma 4.18. Let $\Psi: \Omega \rightarrow \Gamma$ be a monomorphism of finite groups. Let $\bar{\Psi}_k = \Sigma_k \langle \Psi \rangle: \Sigma_k \langle \Omega \rangle \rightarrow \Sigma_k \langle \Gamma \rangle$ ($k \in \mathbb{N}$). Then

$$T_k \Psi_* = (\bar{\Psi}_k)_* T_k.$$

Proof. Similarly to Lemma 4.15, it is easy to show that both compositions are homomorphisms from ‘+’ to ‘ \times ’. Then, it is enough to check the commutativity for Ω -sets, and this can be checked as in [9, Proposition 2.9]. \square

Properties 1 and 4 of Definition 4.4 are valid in $B(S)$ (take for ‘e’ the Σ_1 -set of one element). Then, from Propositions 4.8 and 4.11, we have

Theorem 4.19. The operation composition ‘ \vee ’ defined in Definition 4.5 makes the graded ring $B(S) = \bigoplus_{k \geq 0} B(\Sigma_k)$ an operator ring. \square

Note. Rymer [12] states the same theorem by using the definition of Burnside ring of a compact Lie group to avoid formal negatives in constructing a group from a

semigroup. But he defines the operation $(b \vee a)$ only for elements ' a ' in $B(\Sigma_k)$ (any k) and not for any element ' a ' in $B(S)$.

5. Some notes on β -rings

First we take the definition of β -ring given by Rymer in [12]:

Definition 5.1. A β -ring is a commutative ring C with identity and a composition $B(S) \times C \rightarrow C$ satisfying:

- (1) $(a + b) \cdot c = (a \cdot c) + (b \cdot c)$,
- (2) $(a \times b) \cdot c = (a \cdot c)(b \cdot c)$,
- (3) $(a \vee b) \cdot c = a \cdot (b \cdot c)$,

for all $a, b \in B(S)$, $c \in C$.

Let Γ be a finite group. We define the composition

$$B(\Sigma_r) \times B(\Gamma) \rightarrow B(\Gamma)$$

by means of

$$a \cdot c = (\eta_r)_* \alpha_r^* [\pi_r^*(a) T_r(c)]$$

where

$$\begin{aligned} a &\in B(\Sigma_r), & c &\in B(\Gamma), \\ \alpha_r &: \Sigma_r \times \Gamma \rightarrow \Sigma_r \langle \Gamma \rangle & ((\sigma, g) &\mapsto (g, \dots, g; \sigma)), \\ \pi_r &: \Sigma_r \langle \Gamma \rangle \rightarrow \Sigma_r, & \eta_r &: \Sigma_r \times \Gamma \rightarrow \Gamma. \end{aligned}$$

This composition can be extended by left additivity to all elements in $B(S)$. But it does not seem to be easy to prove that $B(\Gamma)$ becomes a β -ring. In particular, property 3 may be complicated, because it is not enough to take $b \in B(\Sigma_m)$ for some m ; in fact, we have to take $b = b_1 + \dots + b_s$, with $b_j \in B(\Sigma_{m_j})$ ($j = 1, \dots, s$).

Theorem 5.2. The homomorphism $\Psi: B(S) \rightarrow R(S)$ induced by assigning to a finite Σ_r -set X the complex Σ_r -representation $\mathbb{C}(X)$ on the set X as basis is a homomorphism of operator rings.

Proof. The theorem follows from the definitions of ' \vee ' in $B(S)$ and $R(S)$ by showing that Ψ commutes with inductions, restrictions and with the maps T_k ($\bar{\tau}_k$ of [9]). \square

Note that it is not enough to prove $\Psi(b \vee a) = \Psi(b) \vee \Psi(a)$ with $a \in B(\Sigma_r)$ for some $r \in \mathbb{N}$, as Rymer does. In fact, it is not possible to apply [5, Lemma (4.6)] because the set $\{a \in B(\Sigma_r); r \in \mathbb{N}\}$ is not closed under addition.

Let \mathcal{A} be the ring of all λ -operations. By composing $\theta\Psi$, where $\theta: R(S) \rightarrow \mathcal{A}$ is the

map $a \mapsto \tau^a$ (see [9]), we obtain a homomorphism of operator rings $B(S) \rightarrow \mathcal{A}$. So, every λ -ring is a β -ring.

From now on we use the definition of β -ring given by Morris and Wensley [11]:

Let $P = \bigcup_{r \geq 0} \{\Sigma_r\text{-sets}\}$.

Definition 5.3. A β -ring is a commutative ring C with identity together with a function:

$$\beta: P \times C \rightarrow C, \quad \beta(X, c) = \beta(X) \cdot c$$

such that

- (1) $\beta(X + X') \cdot c = \beta(X) \cdot c + \beta(X') \cdot c$,
- (2) $\beta(X \times X') \cdot c = (\beta(X) \cdot c)(\beta(X') \cdot c)$,
- (3) $\beta(X \vee Y) \cdot c = \beta(X) \cdot (\beta(Y) \cdot c)$,
- (4) $\beta(e) \cdot c = c$,

where X, X', Y are elements of P . Both X and X' are Σ_r -sets for the same $r \in \mathbb{N}$. And e is the Σ_1 -set of one element.

A priori Definition 5.3 is weaker than Definition 5.1.

Let Γ be a finite group. We define in $B(\Gamma)$ the β -operations

$$\beta(X) \cdot c = X \cdot c$$

where $X \in P$, $c \in B(\Gamma)$ and $X \cdot c$ is as $a \cdot c$ above.

Rymer proves in [12, Theorem 2] that $B(\Gamma)$ is a β -ring with Definition 5.3, not 5.1. With our definition of the β -operations it is not necessary to use the structure of compact Lie group in order to prove that theorem, because, by [5, Lemma (4.6)], we can suppose c is a Γ -set and then the proof is similar to Rymer's one.

Observe that this answers the question of Boorman [2, p. 149] without using compact Lie groups.

Finally, $B(S)$ is a β -ring (both with Definitions 5.1 and 5.3) with the action

$$\beta(X) \cdot c = X \vee c \quad (X \in P, c \in B(S)).$$

To prove that $B(S)$ is precisely the free β -ring in one generator (with Definition 5.3) is equivalent to show the equivalence between Definitions 5.1 and 5.3. We do not know whether this is true or not; our opinion is that $B(S)$ is, in fact, the free β -ring in one generator.

Acknowledgment

The present work was done during my stay at the Mathematics Institute of Warwick University with the support of a Fleming Award from the British Council. I thank the members of the staff of that institute for their hospitality and, specially, Prof. J.A. Green for his kindness.

My gratitude is also due to the referee for his helpful comments and to Prof. M. Torres, who introduced me to this topic.

References

- [1] M.F. Atiyah and D.O. Tall, Group representations, λ -rings and the J -homomorphism, *Topology* 8 (1969) 253–297.
- [2] E.H. Boorman, S -operations in representation theory, *Trans. Amer. Math. Soc.* 205 (1975) 127–149.
- [3] J.P. Brennan, Products in lambda-rings, University of Illinois at Urbana Champaign, Ph.D., 1984.
- [4] T. tom Dieck, The Burnside ring of a compact Lie group, *Math. Ann.* 215 (1975) 235–250.
- [5] A. Dold, K -theory of non-additive functors of finite degree, *Math. Ann.* 196 (1972) 177–197.
- [6] A. Dress, Notes on the theory of the representations. Part I, *Lecture Notes University of Bielefeld*, 1970.
- [7] A. Dress, Operations in representation rings, *Proc. Sympos. Pure Math.* 21 (1971) 39–45.
- [8] J.A. Green, Axiomatic representation theory, *Proc. Sympos. Pure Math.* 21 (1971) 57–64.
- [9] P. Hoffman, τ -rings and Wreath Product Representations, *Lecture Notes in Mathematics* 746 (Springer, Berlin, 1979).
- [10] D. Knutson, λ -rings and the Representation Theory of the Symmetric Group, *Lecture Notes in Mathematics* 308 (Springer, Berlin 1973).
- [11] I. Morris and C.B. Wensley, Adams operations and λ -operations in β -rings, *Discrete Math.* 50 (1984) 253–270.
- [12] N.W. Rymer, Power operations on the Burnside ring, *J. London Math. Soc.* (2) 15 (1977) 75–80.